

EVALUATION OF THE EFFICIENCY OF EFFECTIVE MICROORGANISM PREPARATION EM1 IN RESISTING SOME FUNGI THAT CAUSE TOMATO ROOT ROT DISEASE

¹Shahla abdukkareem sattar and ¹Ahed A. H. Matloob*

¹ Al-Mussaib Technical College, University of Al-Furat Al-Awsat Technical, 51009, Babylon, Iraq.

Corresponding author: Shahla.abdukkareem.tcm.9@student.atu.edu.iq

Received (15/09/2025), Received in revised form (25/10/2025)

Accepted (05/11/2025), Available online (31/12/2025)

FJIAS 2025, 1(4): 26-34

Abstract: *This study aimed to isolate some pathogenic fungi from tomato plants infected with root rot disease, diagnose them, test their pathogenicity, and try to control the causes of tomato root rot disease using EM-1 microorganism preparations under laboratory conditions. The results of the field survey that was conducted in some fields of Karbala, Baghdad and Babylon governorates showed the spread of root rot disease in the areas covered by the field survey, with an infection rate ranging between 62-100% and an infection severity of 35-57%. Several types of fungi were isolated and identified from the roots of tomato plants infected with root rot disease. The most frequent pathogenic fungi were *Fusarium solani*, followed by *Rhizoctonia solani* and *Macrophomina phaseolina*. The results showed that all tested fungi isolates were pathogenic and led to any significant reduction in the percentage of germination, compared to the comparison treatment. The results showed the ability of the EM1 microorganism preparation to inhibit the growth of pathogenic fungi *F.solani* Fs1, *R.solani* Rs5 and *M. phaseolina* Mp8 on PDA culture medium. The EM1 showed the highest effect at 5, 10 and 15% concentrations on pathogenic fungi growth, as the percentage reached The percentage of inhibition is 100%.*

Keywords: Tomato, fungi, root rot, Effective microorganism, disease.

1. INTRODUCTION

The tomato crop is affected by a group of pathogens that have different pathogenic mechanisms that reduce the yield of tomato and affect the quality of the product [1] and [2]., and many fungi, bacterial species, viruses and some types of Mycoplasma are included under the pathogens [3] and among these important diseases are *Fusarium* wilt, leaf rot, root knot, early blight on tomatoes and others fungi and the most famous of them are *Fusarium solani*, *Rhizoctonia solani*, *Microphomina phaseolina*, *Fusarium oxysporium* and *Pythium* spp. [4] and [2]. Excessive use of chemical pesticides in combating diseases led to many environmental problems and toxic effects for humans and non-target organisms, which motivated researchers to develop alternative inputs to combat plant diseases, including the use of biological control [5], biological control agents act as alternatives For pesticides, these organisms are used as biocides and fertilizers in sustaining and enhancing crop production and protection and are usually biological strains isolated from the soil [6] and [7]. Induced resistance was also used that works on Stimulating plants and pushing them to resist disease, whether through the use of biological or non-biotic product Its production

promised a good alternative to the control of plant pathogens, and the application of Nanotechnology in agriculture showed promising results in terms of Related to the development and use of products such as fertilizers and pesticides that significantly and significantly enhanced the induction of resistance in plants [8], [9] and [10]. The biologically safe techniques based on effective microorganisms (EM1 Effective Micro-Organisms) were used to maintain soil fertility, many of which are highly resistant to toxic substances and have positive effects on plant growth and plant protection from toxic substances under different stress conditions [11]. Because of the importance of the disease of seedling death and tomato root rot and to search for methods of resistance in line with modern global trends in reducing the use of chemicals in agriculture, this study aimed to isolate some pathogenic fungi from tomato plants infected with root rot disease, diagnose them, test their pathogenicity and try to combat the causes of root rot disease. Tomato using some normal and Nano chemical inducing agents and EM-microorganism preparation under laboratory conditions.

2. MATERIALS AND METHODS

2- 1. field survey

A field survey was conducted for the season 2022-2021 in three governorates of Baghdad, Babylon and Karbala for the period 20/6/2021 to 5/9/2021. Samples that showed symptoms of root rot disease represented by wilting, yellowing of leaves and general weakness in growth were collected, noting the presence of rots. It has a brown color on the main and subsidiary roots by removing the plant randomly and located within the intersection of the diameters for each site. The healthy and infected plants were examined and then placed in bags (polyethylene) and marked and kept in the refrigerator at a temperature of 4°C until the fungi accompanying the roots were isolated from each sample. The percentage of infection was calculated and The percentage of infection severity for the root system was calculated according to the pathological index consisting of 6 degrees, as follows :0=healthy plant, 1 = More than 0–25% of the roots are affected, 2 = More than 25%-25% of the roots are infected 3=More than 75%-50% of the roots are infected, 4= more than 75% of the roots are infected, 5 = plant death

in a dark color or the death of the plant. The percentage of severity was calculated according to the Mckinney equation [12] as follows:

Severity (%) = ((Plants in 1 degree ×1+... Plants in 5 degree×5)/ all plants ×5) ×100%.

2-2 Isolation and identification of fungi associated the infected roots of tomato plants

The process of isolation was carried out from the tomato plant samples that showed symptoms of infection on the next day of the field survey, as the plants were brought to the laboratory and the affected roots were washed with running water to remove dust for 5 minutes, and they were cut into small pieces with a length of 0.5 cm, and superficially sterilized by dipping them with a solution of Sodium hypochlorite (1% free chlorine) for one minute, then washed with sterile distilled water for two minutes and dried with sterile filter paper. Potato Dextrose Agar (PDA) sterilized by an autoclave device at a temperature of 121°C and a pressure of 1 atmosphere for 15-20 minutes, and the antibiotic Tetracycline was added to it. Small pieces from the ends of the fungal hyphae were placed in the center of a petri dish containing PDA medium. The dishes were incubated for 4 days in the incubator and examined under the compound microscope. The fungi were diagnosed for the level of sex and species after the appearance of the fungal growths and depending on the characteristics of the colony. Form and the nature of the mycelium, the spores and the structures that they form using the approved taxonomic keys [13, 14, 15,16]. by Dr. Ahed Abd Ali Hadi. The appearance of fungi was calculated as following: repeat the fungus in the sample = (The number of fungus appeared in dishes\Total number of pieces used in the sample)x 100.

2-3 Detection of pathogenic isolates of *Rhizoctonia solani*, *Fusarium solani*, and *Macrophomina phaseolina* using cabbage seeds on Potato Dextrose Agar (PDA) medium.

Seven isolates of *M. phaseolina*, five isolates of *F. solani* and 10 isolates of *R. solani* were tested. According to the method of Bolkan and Butler [17], Petri dishes with a diameter of 9 cm were prepared containing PDA sterilized medium with an autoclave for 20 minutes and the antibiotic was added Bio-tetracycline to the medium, the plates were inoculated from their center after hardening the medium with a 0.5 cm disc taken from the edges of the colony of the fungi *F. solani*, *R. solani* and *M. phaseolina* at the age of five days, then the plates were incubated in the incubator at a temperature of 25 ± 1 °C for three days, after that, the seeds were used after superficial sterilization with sodium hypochlorite solution (free chlorine 1%), which was previously tested for germination capacity in a circular motion near the edge of the plate, at a rate of 10 seeds per plate, with three replicates for each isolate as well as a control treatment (without pathogenic fungi). The dishes were incubated in the incubator at a temperature of 25 ± 1 °C. After seven days, the results were taken by calculating the percentage of germination.

2-4 Evaluation of the antagonistic ability of effective microorganisms EM1 to inhibit the growth of *Rhizoctonia solani*, *Fusarium solani* and *Macrophomina phaseolina* on PDA medium.

The inert basic EM1 solution, were activated by mixing it with molasses, chlorine-free warm water at a ratio of 5 ml of inert EM1 solution, 5 ml of molasses, and 95 ml of chlorine-free water. A warm place away from sunlight for 20 days at a temperature between 35-40 C. During this period the package was opened more than once to leak gas and form a layer of precipitate at the bottom of the package. Add 1, 3, 5, 10 and 15% ml of activated EM1 solution to a beaker. 100 ml glass was completed and the volume was completed to 100 ml of PDA culture medium, each concentration individually, then the mixture was poured into Petri dishes. The dishes was inoculated by isolates of *F. solani* (Fs1) and *R. solani* (Rs5) and *M. phaseolina* (Mp8) at 5 days old in the center of the dish with 3 replicates, the plates were placed in the incubator at a temperature of $25 + 1$ °C for 7 days and the colony diameter and the percentage of inhibition were measured based on the comparison treatment to evaluate the efficiency of the EM1 preparation according to the following equation:

$$\% \text{ Inhibition} = [(R - r) / R] \times 100.$$

Where, r is the radius of the fungal colony against the bioagents and R is the radius of the fungal colony without the bioagents.

3- RESULTS AND DISCUSSION:

3-1 Field survey of tomato root rot

The results of Table 1 showed the field survey that was conducted in some fields of Karbala, Baghdad and Babylon governorates, the spread of root rot disease in the areas covered by the field survey, with an infection rate ranging between 62-100% and an infection severity of 35-57%. Al-Azzawiya and Al-Haidaria, which reached 80, 85, 85, 96 and 100%, respectively. The reason for the high infection rate in these fields may be attributed to the repeated cultivation of the tomato crop in them or the cultivation of other crops belonging to the Solanaceae family in the same fields, which leads to the accumulation of fungal pollen for pathogens, especially sclerotia, which remain in the soil for a long time, and appropriate environmental conditions, especially temperatures suitable for the growth of pathogenic fungi, hoeing and weeding processes, lead to the making of wounds in the roots, which facilitates the process of fungal invasion and infection [18]. While the highest severity of infection was in the fields of Tarmiyah, which amounted to 57%, followed by Al-Azzawiya area with 54%, and then Al-Haidariya area with 57%. While the infection severity was less in the Bada'a and Al Mahaweel region, reaching 35%, and the reason for this is that the field was newly planted with the tomato crop, and the severity of the infection varied depending on the operations of crop service, soil, regular irrigation and control methods, as well as the

difference in environmental factors such as temperature and humidity, which have a significant impact on Increase in fungal infection.

Table 1. Percentage of infection and severity of tomato root rot disease for some regions for the agricultural season 2021-2022

Region	Disease induce (%)	Severity (%)
Babel / Al-Azzawiya	96	54
Babylon / Tahiriya	62	42
Babylon / heresy	85	35
Babylon / Mahaweel	77	35
Baghdad / Tarmiyah	85	47
Baghdad / Yusufiyah	66	43
Baghdad / Latifiya	80	42
Karbala / Haidaria	100	57
Karbala / Khan Al-Nass	71	40
Karbala / Husseiniya	70	44

3-2 Isolation and identification of fungi accompanying the roots of infected tomato plants

Several types of fungi were isolated and diagnosed from the roots of tomato plants infected with root rot disease, (Table 2) and the most frequent pathogenic fungus was *Fusarium solani*, which was isolated from all areas covered by the survey, its appearance rate was 55.9% and the highest percentage of its appearance was 91%, followed by the fungus *Rhizoctonia solani*, which was isolated from seven regions, at a rate of 61.8%, and the highest rate of its appearance was 95%, followed by the fungus *Macrophomina phaseolina*, which was also isolated from seven regions, where the highest percentage of its appearance was 92%, with a rate of 58.5%. These results show that the fungi *F.solani* , *R.solani* and *M. phaseolina* are among the most important pathogens associated with root rot that infect many field crops.

Table 2. Percentage of the appearance of fungi in the roots of tomato plants infected with root rot.

Fungus	Appearance rate (%)	Highest ratio of appearance
<i>Fusarium solani</i>	55.9	91
<i>Rhizoctonia solani</i>	61.8	95
<i>Macrophomina phaseolina</i>	58.5	92
<i>Pencillium spp.</i>	52.6	81
<i>Aspergillus nigeria</i>	29.16	73
<i>Chaetomium globosum</i>	32.33	44
<i>Acremonium spp.</i>	13	13
<i>Lichtheimia ramosa</i>	16	16

These results are consistent with many studies that confirmed the association of these fungi with root rot diseases in many crops [19]. The diagnostic results showed the presence of many Fungi associated the roots of tomatoes and less frequent, such as *Pencillium* spp, *Aspergillus niger*, *Chaetomium globosum* and *Acremonium* spp. and *Lichtheimia ramosa* with a higher percentage of 81, 73, 44, 13 and 16%, respectively. Studies agree with what was found by Ismael and Salih [20], Alsudani, Al-awsu [21], Muslim et al, [22]. that these fungi *F. solani* , *R. solani* and *M. phaseolina* infect tomato and cause root rot and seedling fall diseases.

3.3 Testing the pathogenicity of isolates of pathogenic fungi using cabbage seeds on PDA . culture medium

The results of Table 3 showed that all tested fungi isolates led to any significant reduction in the percentage of germination, compared to the comparison treatment in which the percentage of seed germination was 100%.

Table 3. Detection of the pathogenic isolates of *Fusarium solani*, *Rhizoctonia solani* and *Macrophomina phaseolina* using cabbage seeds on PDA medium.

Region	Isolate symbol	No. germinated seeds	Germination (%)
Babel / Al-Azzawiya	control	10.00	100.00
Babylon / Tahiriya	Fs1	1.33	13.33
Babylon / heresy	Fs2	3.67	36.67
Babylon / Al-Mahaweel	Fs3	2.67	26.67
Baghdad / Tarmiyah	Fs4	5.00	50.00
Baghdad / Yusufiya	Fs5	3.33	33.33
Baghdad / Latifiya	Fs6	2.67	26.67
Karbala / Haidari	Fs7	4.33	43.33
Karbala / Khan Al-Nass	Fs8	2.67	26.67
Karbala / Hussainiya	Fs9	6.67	66.67
Babylon / heresy	Fs10	4.33	43.33
Baghdad / Tarmiyah	Rh3	2.33	23.33
Baghdad / Yusufiya	Rh5	0.00	0.00
Karbala / Haidari	Rh6	4.67	46.67
Karbala / Hussainiya	Rh8	1.67	16.67
Babel / Al-Azzawiya	Rh10	6.33	63.33
Babylon / Tahiriya	Mp1	5.00	50.00
Babylon / heresy	Mp2	4.67	46.67
Babel Al / Mahaweel	Mp3	6.00	60.00
Baghdad / Tarmiyah	Mp4	3.33	33.33
Karbala / Haidari	Mp5	3.00	30.00
Karbala / Hussainiya	Mp8	0.33	3.33
LSD (0.05)	-	0.815	8.16

Each number in the table represents an average of three replicates, Fs = *Fusarium solani* , Rs = *Rhizoctonia solani* , Mp = *Macrophomina phaseolina* The number near the isolate symbol represents the isolate number

The isolate of the fungus *F.solani* (Fs1) excelled in Babil / Al-Azzawiya governorate in reducing the percentage of germination, as the average percentage of germination in it was 13% followed by isolates of Fs3, Fs6 and Fs8, where the percentage of germination reached 26.67%, while the percentage of germination in the rest of the isolates ranged between 36.67 - 66.67%, as the results of the table 3 indicated that all isolates The tested fungus *R.solani* caused a significant reduction in the germination of the seeds of cabbage compared to the comparison treatment, and the isolates varied among themselves in reducing the germination rate. Rs8 Isolation of Karbala / Al-Haidariya governorate, in which the percentage of seed germination reached 16.67%, while the percentage of germination in the rest of the isolates ranged between 23.33 - 63.33%, and the results in the table 3 showed that all isolates of *M. phaseolina* showed a reduction Significant in the percentage of germination of cabbage seeds compared to the comparison treatment, which reached the percentage of germination 100%, and it was more superior among isolates of Mp8 isolate belonging to Karbala / Al-Haidariya governorate, the percentage of germination in it was 3.33%, followed by isolate Mp9 isolate Karbala Al-Hussainiya governorate and the percentage of germination was 23.33%, while the percentage ranged For the rest of the isolates between 30.00-60.00% .

The reason for the variation of isolates in their effect on the percentage of germination of cabbage seeds is due to genetic differences between isolates of fungi collected from several regions in different governorates, or the difference of these isolates in their ability to produce toxins or enzymes degrading pectin and cellulose in the early stages of infection and these enzymes Phosphatase, pectinase, cellulase, pectin methylesterase, pectin lyase, which have an effective role in the pathogenesis of fungi, have a significant role in penetrating the plant host and causing infection. This was confirmed by Wrather [23], In this test, the isolates with the most reduced germination of cabbage seeds, namely Fs1, Rs5 and Mp8, were selected for subsequent tests.

3-4 Evaluation of the antagonistic ability of effective microorganisms (EM1) in inhibiting the growth of the fungi *Rhizoctonia solani*, *Fusarium solani* and *Macrophomina phaseolina* on PDA medium.

The test results are shown in Table 4. The ability of the EM1 microorganism preparation to inhibit the growth of pathogenic fungi *F.solani* Fs1, *R.solani* Rs5 and *M. phaseolina* Mp8 on PDA culture media. The microorganism EM1 showed the highest effect at 5, 10 and 15% concentrations on the growth of pathogenic fungi, as the percentage of inhibition reached 100% compared with the control treatment of fungi alone, where the percentage of inhibition was 0.00%. While the concentration of 1 and 3 ml showed a decrease in the growth level of the colony diameter of the fungus Fs1, which reached 2.55 and 1.33 cm, respectively, and the percentage of inhibition was 71.29 and 81.18%, respectively. As for the isolate of the fungus *R.solani* Rs5, the diameter of the colony growth of the fungus at concentrations 1 and 3% was 2.67 and 1.25 It was clear that the higher the concentration, the higher the percentage of inhibition, reaching 70.37 and 86.11%, respectively, and the isolate of *M. phaseolina* Mp8. The percentage of inhibition at concentration 1 and 3 ml was 83.33 and 87.96%, respectively, and the diameter of the mushroom colony growth was 1.50 and 1.08. Straight. The results showed that there is a direct proportion to the percentage of inhibition with an increase in the concentration of the biological preparation EM1 if it causes a significant reduction in the growth of pathogenic fungi, and the greater the concentration of the preparation, the higher the percentage of inhibition compared to the treatment of fungi alone, which amounted to 0.00%.

The inhibition of the pathogenic fungus is successful due to the by-products produced by microorganisms such as the antibiotics produced by the bacteria present in the EM1 antibiotic during fermentation, which directly affects the growth of many pathogens as well as the presence of yeasts and fungi [24].

Table 4. Effect of EM1 on the growth of the pathogenic fungi *Rhizoctonia solani*, *Fusarium solani* and *Macrophomina phaseolina* on PDA medium.

Treatment	Concentration	Colony diameter.cm	Inhibition (%)
control	0	9.00	0.00
EM1 + fs1	1	2.58	71.29
	3	1.33	85.18
	5	0.00	100.00
	10	0.00	100.00
	15	0.00	100.00
control	0	9.00	0.00
EM1 + Rs5	1	2.67	70.37
	3	1.25	86.11
	5	0.00	100.00
	10	0.00	100.00
	15	0.00	100.00
control	0	9.00	0.00
EM1 + Mp8	1	1.50	83.33
	3	1.08	87.96
	5	0.00	100.00
	10	0.00	100.00
	15	0.00	100.00
LSD (0.05)	-	0.095	1.047

Each number represents an average of three replicates Fs1 = *Fusarium solani* , Rs5 = *Rhizoctonia solani* , Mp8 = *Macrophomina phaseolina*, the number near the symbol represents the isolate number

4-CONCLUSIONS

We concluded from the current study the spread of tomato root rot disease in all areas covered by the field survey. The pathogenic fungi *Fusarium solani*, *Rhizoctonia solani* and *Macrophomina phaseolina* were found to be the main causes of tomato root rot diseases in Baghdad, Babylon and Karbala governorates. Ordinary the microorganism preparation had high antagonistic ability against root rot fungi *F. solani*, *R. solani* and *M. phaseolina* under laboratory and field conditions.

REFERENCES

- [1] Al-Ani, L.K.T.; Furtado, E.L.2020. The effect of incompatible plant pathogens on the host plant. In Molecular Aspects of Plant Beneficial Microbes in Agriculture; Academic Press: Cambridge, MA, USA,; pp. 47–57. Ebert, A. W. 2020. The role of vegetable genetic resources in nutrition security and vegetable breeding. *Plants*, 9(6), 736.
- [2] Campos, M. D., Félix, M. D. R., Patanita, M., Materatski, P., Albuquerque, A., Ribeiro, J. A., & Varanda, C. 2022. Defense strategies: The role of transcription factors in tomato–pathogen interaction. *Biology*, 11(2), 235.
- [3] Campos, M. D., Patanita, M., Campos, C., Materatski, P., Varanda, C. M., Brito, I., & Félix, M. D. R. 2019. Detection and Quantification of *Fusarium* spp.(*F. oxysporum*, *F. verticillioides*, *F. graminearum*) and *Magnaportheiopsis maydis* in Maize Using Real-Time PCR Targeting the ITS Region. *Agronomy*, 9(2), 45. Manisha, M., Arpita, S., & Tomar, B. S. 2018. Genetic and molecular regulation of colour and pungency in hot pepper (*Capsicum* spp): a review. *Indian Journal of Agricultural Sciences*, 88(3), 343-353.
- [4] Agrios, G.N. .2005. plant pathology, 5th edn. Academic Press, New York, p:922.
- [5] Aboutorabi, S. N., Nasiriboroumand, M., Mohammadi, P., Sheibani, H., & Barani, H. 2018. Biosynthesis of silver nanoparticles using safflower flower: structural characterization, and its antibacterial activity on applied wool fabric. *Journal of Inorganic and Organometallic Polymers and Materials*, 28(6), 2525-2532.
- [6] Fasusi, O. A., Cruz, C., & Babalola, O. O. 2021. Agricultural sustainability: microbial biofertilizers in rhizosphere management . *Agriculture*, 11(2), 163.
- [7] Elnahal, A. S., El-Saadony, M. T., Saad, A. M., Desoky, E. S. M., El-Tahan, A. M., Rady, M. M., ... & El-Tarabily, K. A. 2022. The use of microbial inoculants for biological control, plant growth promotion, and sustainable agriculture: A review. *European Journal of Plant Pathology*, 1-34.
- [8] Kah, M., Tufenkji, N., & White, J. C. 2019. Nano-enabled strategies to enhance crop nutrition and protection. *Nature nanotechnology*, 14(6), 532-540.
- [9] Lowry, G. V., Avellan, A., & Gilbertson, L. M. 2019. Opportunities and challenges for nanotechnology in the agri-tech revolution. *Nature nanotechnology*, 14(6), 517-522.
- [10] Paramo, L. A., Feregrino-Pérez, A. A., Guevara, R., Mendoza, S., & Esquivel, K. 2020. Nanoparticles in agroindustry: Applications , toxicity, challenges, and trends . *Nanomaterials* , 10(9), 1654.
- [11] Ivanova, S. V., & Ryabchikova, I. A. 2022. Application of Microbiological Preparations in the Cultivation of Agricultural Crops with the Aim to Reduce the Technogenic Load of Metallurgical Plants. In IOP Conference Series: Earth and Environmental Science (Vol. 988, No. 3, p. 032060). IOP Publishing.
- [12] Mckinney, H.H. 1923. Influence of soil temperature and moisture on infection of wheat seedling by *Helminthosporium sativum*. *J. Agric. Research* 26: 195–217.
- [13] Parmeter, J. R. and H. S. Whitney. 1970. Taxonomy and nomenclature of the imperfect stage In: *Rhizoctonia solani* Biology and pathology. Parmeter, J. R. Univ. of California . 7–19.

- [14] Ellis, M.B. 1971. Dematiaceous Hyphomycetes. Commonwealth Mycological Institute Kew , survey England .1- 608.
- [15] Booth, C. 1977. Fusarium laboratory guide to the identification of the major species. Commonwealth Mycological Institute Kew, Surrey, England. 58 pp.
- [16] Summerell , B . A . and Leslie , J. F. 2006. The Fusarium Laborator manual. 388pp.
- [17] Bolkan, H. H., and E. E. Butler . 1974. Studies on heterokaryosis virulence of *Rhizoctonia solani* . Phytopathology 64: 513 – 522.
- [18] El-Mougy, N. S., M. D. I. H. Aly, E. I. Imbabi and M. M. Abdel-Kader. 2011. First Record of Sclerotinia Foliage Blight Disease on Pepper under Protected Cultivation System in Egypt.
- [19] Emmanuel, A., Dawar, S., and Zaki, M. J. 2010. Effect sida pakistanica s. Abedin and senna holosericea fresen on growth and root rot diseases of okra and mash bean. Pak. J. Bot, 42(1), 391-400.
- [20] Ismael, J. H., & Salih, R. M. 2016. Integrated Management of Tomato Damping-off Disease in Bekrajo, Sulaimani Governorate, Kurdistan Region, Iraq. Jordan Journal of Agricultural Sciences, 12(1).
- [21] Alsudani, A. A. 2020. Biocontrol of *Rhizoctonia solani* (Kühn) and *Fusarium solani* (Marti) causing damping-off disease in tomato with *Azotobacter chroococcum* and *Pseudomonas fluorescens*. Pakistan Journal of Biological Sciences: PJBS, 23(11), 1456-1461.
- [22] Muslim, S. N., Aziz, R. A. R., & Al-Hakeem, A. M. (2021). Biological control of *Azotobacter chroococcum* on *Fusarium solani* in tomato plant. In Journal of Physics: Conference Series (Vol. 1879, No. 2, p. 022018). IOP Publishing.
- [23] Wrather, J. A.; Anderson, T. R., Arsyad, D. M., Gai, J., Ploper, L. D.; Porta- Puglia , A.; Ram, H. H. and Yorinori, J. T. 1997. Soybean disease lossestimates for the top 10 soybean producing countries in 1994. Plant Dis. 81: 107-110 .
- [24] Higha, T. (2006). An Earth Saving Revolution. (English Translation), Sunmark Publishers, Inc. Tokyo, Japan.